

Delaro®

Net contents : 5 L, 15 L

Fungicide

Claim statement: systemic fungicide for the control of various fungal diseases in cereals


Contains:

175 g/litre prothioconazole and 150 g/litre trifloxystrobin in the form of an suspension concentrate

Registered pursuant to the ACVM Act 1997, No P8953
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, No HSR100886
See www.epa.govt.nz for approval conditions

DG Information



UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S
(Trifloxystrobin, Prothioconazole Solution)
PACKING GROUP III
CLASS 9
HAZCHEM CODE 3Z

IN A TRANSPORT EMERGENCY DIAL 111 POLICE OR FIRE BRIGADE



Barry

WARNING
KEEP OUT OF REACH OF CHILDREN
HSNO Classifications:



May cause damage to organs through prolonged or repeated exposure
Very toxic to aquatic life with long lasting effects

PRECAUTIONS:

Do not eat, drink or smoke while using this product.

Avoid breathing dust/mist/vapours/spray. Use only outdoors or in a well –ventilated area.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

Do not apply directly into or onto water

PERSONAL PROTECTION:

When handling, wear full protective clothing such as impervious gloves, respirator, waterproof hat, coat and trousers (worn outside rubber boots).

STORAGE: Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled, move the victim to fresh air immediately.



Barayan

It is an offence to use this product on animals.

DIRECTIONS FOR USE:

MIXING: Shake container well before use. Half fill the spray tank with water, then pour in the required quantity of Delaro® with agitators running, then top up with water. Do not pre-mix.

RESISTANCE MANAGEMENT

Group	3 + 11	Fungicide
-------	---------------	-----------

Delaro contains prothioconazole from the DMI (DeMethylation Inhibitor) mode of action (Group 3) and trifloxystrobin from the Quinone outside Inhibitors (QoI) mode of action (Group 11). Resistance to these fungicides and related products could develop in some disease situations from repeated use. To minimize this risk use a maximum of 2 applications per season. Make no more than 2 applications using fungicides from the QoI group and no more than 3 applications, including seed treatment, from the DMI group.

RECOMMENDATION:

Crop	Disease	Product Rates	Comments
Wheat	Stripe rust Leaf rust Speckled leaf blotch Glume blotch	600 – 750 ml/hectare	Apply on first appearance of disease. Use the higher rate when disease pressure is high. Should disease pressure persist or re-infection occur, a further application may be required 3-4 weeks later.
Barley	Scald Net blotch Ramularia leaf spot Leaf rust	600 – 750 ml/hectare	Apply on first appearance of disease. Use the higher rate when disease pressure is high. Should disease pressure persist or re-infection occur, a further application may be required 3-4 weeks later.
Ryegrass	Crown rust Stem rust Blind seed disease	600 – 750 ml/hectare	Apply on first appearance of disease. Use the higher rate when disease pressure is high. Should disease pressure persist or re-infection occur, a further application may be required 3-4 weeks later.

Apply Delaro in at least 200 litres of water per hectare by ground or 50 litres per hectare by aerial application.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

WITHHOLDING PERIOD:

Wheat, barley (grain) - 56 days
Wheat, barley (forage) - 42 days
Ryegrass (forage) - 35 days
Ryegrass (seed) - 35 days



CONDITIONS OF SUPPLY: READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Date of Manufacture:

Batch No:

Delaro® is a registered trademark of the Bayer Group

Bayer

Registered to:

Bayer New Zealand Limited

B:HIVE Building, 74 Taharoto Rd, Smales Farm

Takapuna, Auckland, 0622, New Zealand

Ph. 0800 428 246

EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607

