

# Escolta®

Net contents : 5 L

## ***Fungicide***


A fungicide for the control of foliar diseases in fodder and sugar beets

Contains: 160g/litre cyproconazole and 375 g/litre trifloxystrobin in the form of a suspension concentrate

Registered pursuant to the ACVM Act 1997, No P9302  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to the HSNO Act 1996, No HSR101050  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

### DG Information



UN 3082  
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID,  
N.O.S. (TRIFLOXYSTROBIN, CYPROCONAZOLE),  
PACKING GROUP III  
CLASS 9  
HAZCHEM CODE 3Z

IN A TRANSPORT EMERGENCY DIAL 111 POLICE OR FIRE BRIGADE



03-07-2017

**WARNING****KEEP OUT OF REACH OF CHILDREN**

**Approved Handler:** This product must be under control of an approved handler when applied in a wide dispersive manner or when used by a commercial contractor

**HSNO Classifications:**

6.1E May be harmful.

6.8B Suspected of damaging fertility or the unborn child

6.9B May cause damage to organs through prolonged or repeated exposure

9.1A Very toxic to aquatic life with long lasting effects

**PRECAUTIONS:**

Do not eat, drink or smoke while using this product.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Avoid contact with the skin and eyes

**PERSONAL PROTECTION:**

When handling, wear full protective clothing such as impervious gloves, respirator, waterproof hat, coat and trousers (worn outside rubber boots).

**STORAGE:** Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

**SPILLS AND DISPOSAL:** When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

**Disposal:** Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

**Container Disposal:** Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

**FIRST AID:**

If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled, move the victim to fresh air immediately.



03-07-2017

**It is an offence to use this product on animals.**

**DIRECTIONS FOR USE:**

**MIXING:** Shake container well before use. Half fill the spray tank with water, then pour in the required quantity of Escolta with agitators running, then top up with water. Do not pre-mix.

**RESISTANCE MANAGEMENT**

**Group 3 + 11 Fungicide**

Escolta contains trifloxystrobin from the strobilurin mode of action (Group 11) and cyproconazole from the DMI (DeMethylation Inhibitor) mode of action (Group 3). Resistance to these fungicides and related products could develop in some disease situations from repeated use. To minimize this risk use a maximum of 2 applications per season.

**RECOMMENDATION:**

Crop	Disease	Product Rate	Water rate	Remarks
Fodder Beet Sugar Beet	Rust Powdery mildew Cercospora Ramularia leaf spot	0.35 L/ha	Ground: Apply at least 200 litres of water per hectare  Aerial: 80 L/ha	Apply at the onset of disease. Escolta will give most effective disease control when applied as the diseases first becomes active in the crop and before disease becomes established. It will give prolonged protection from re-infection, but a second spray may be applied 3-4 weeks later if crops are at risk from later attacks.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**WITHHOLDING PERIOD:**

Fodder Beet/Sugar Beet: 42 days

**CONDITIONS OF SUPPLY: READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.**

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Date of Expiry:

Batch No:

June/2017



03-07-2017

TNP® is a registered trademark of the Bayer Group

**Bayer**

Registered to:  
Bayer New Zealand Limited  
3 Argus Place, Hillcrest, Auckland 0627

Ph. 0800 428 246  
EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607



*[Handwritten signature]*  
03-07-2017