

Prosaro®

Net contents : 5 L, 15 L

Fungicide


Claim statement: systemic fungicide for the control of diseases in wheat, barley and ryegrass seed crops.

Contains: 125 g/l prothioconazole and 125 g/l tebuconazole in the form of an emulsifiable concentrate. Also contains > 10% N,N-dimethyl decanamide.

Registered pursuant to the ACVM Act 1997, No P7662
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, No HSR007871
See www.epa.govt.nz for approval conditions

DG Information



UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID. N.O.S.
(TEBUCONAZOLE, PROTHIOCONAZOLE SOLUTION)
PACKING GROUP III
CLASS 9
HAZCHEM CODE 3Z

IN A TRANSPORT EMERGENCY DIAL 111 POLICE OR FIRE BRIGADE



03/03/2021

WARNING
KEEP OUT OF REACH OF CHILDREN
HSNO Classifications:



- 6.1E May be harmful if swallowed or inhaled.
- 6.3B Causes mild skin irritation.
- 6.4A Causes serious eye irritation.
- 6.9B May cause damage to organs through prolonged or repeated exposure.
- 9.1B Toxic to aquatic life with long lasting effects.

PRECAUTIONS:

- Read label before use.
- Do not eat, drink or smoke while using this product.
- Do not breathe dust/fume/mist/vapours/spray.
- Remove protective clothing and wash hands and face thoroughly before meals and after work.
- Avoid contact with the skin and eyes.
- Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area
- Do not apply directly into or onto water

PERSONAL PROTECTION:

When handling, wear full protective clothing such as impervious gloves, respirator, waterproof hat, coat and trousers (worn outside rubber boots).

STORAGE: Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled, move the victim to fresh air immediately.



03/03/2021

It is an offence to use this product on animals.

DIRECTIONS FOR USE:

MIXING: Half fill the spray tank with water, then pour in the required quantity of **Prosaro**[®] with agitators running, then top up with water. Do not pre-mix.

RESISTANCE MANAGEMENT

Group	3	Fungicide
-------	----------	-----------

Prosaro[®] is a member of the DMI group of fungicides. Resistance to this fungicide and related DMI products could develop in some disease situations from repeated use. To minimize this risk, use strictly in accordance with label instructions. In cereal and ryegrass seed crops, use a maximum of 3 DMI applications, including the seed treatment per crop at full recommended label rates.

RECOMMENDATION:

Crop	Disease	Product Rates	Comments
Wheat	Glume blotch Leaf rust Powdery mildew Speckled leaf blotch Stripe rust	1 L/ha	Apply on first appearance of disease. Should disease pressure persist or reinfection occurs, a further application may be required 3 – 4 weeks later.
	Ear disease complex		Apply to fully-emerged ears.
Barley	Leaf rust Net blotch Ramularia leaf spot Scald Spot blotch	1 L/ha	Apply on first appearance of disease. Should disease pressure persist or reinfection occurs, a further application may be required 3 – 4 weeks later.
	Ear disease complex		Apply to fully-emerged ears.
Ryegrass Seed Crops	Crown rust Stem rust	1 L/ha	Apply on first appearance of disease. Should disease pressure persist or reinfection occurs, a further application may be required 3 – 4 weeks later.
	Blind seed disease		Apply at start of flowering and a second application 10 – 14 days later.

Apply **Prosaro**[®] in 200 litres of water per hectare by ground-based methods or 50 litres per hectare by aerial application. Use high water volumes in dense or advanced crops especially under dry weather conditions. **Prosaro**[®] may cause yellow leaf spotting when applied after head emergence to some durum wheats, especially Tara, and yellowing when applied with emulsifiable concentrate herbicide formulations in barley.

COMPATIBILITY

Prosaro is compatible with commonly-used herbicides and insecticides.



It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD:

Cereals (grain) - 56 days
Cereals (forage) - 42 days
Ryegrass (seed) - 30 days
Ryegrass (forage) - 35 days

CONDITIONS OF SUPPLY: READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Date of Expiry:

Batch No:

Prosaro® is a registered trademark of the Bayer Group

Bayer

Registered to:

Bayer New Zealand Limited

B:HIVE Building, 74 Taharoto Rd, Smales Farm

Takapuna, Auckland, 0622, New Zealand

Ph. 0800 428 246

EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607



A handwritten signature in blue ink, appearing to be "John G. ...".

03/03/2021