

Updated 2015

BAYER UTE CARD™

Convenient “4 in 1” herbicide for post emergence weed control in Fodder and Red Beet



Betanal[®]
quattro

Benefits

Betanal quattro is a “4 in 1” formulation, containing the four key active ingredients used in herbicides for beet crops in one convenient product.

Betanal quattro provides ease of use, reduces the chances of mixing errors and reduces tank filling time.

Betanal quattro lessens packaging, including storage, transport and disposal.

Betanal quattro is hard on weeds but has a high level of safety to the crop.

Betanal quattro provides flexibility as it can be used from the beet cotyledon stage onwards.



Scan the QR Code to access further Betanal quattro technical information



| | |
|--------------------------|---|
| Rate | 2-5 L/ha |
| Spray Timing | Use 2 L/ha from the cotyledon to 1st true leaf stage. Use 3-5 L/ha after the crop has 2 true leaves fully developed. |
| Weed size | Apply to actively growing weeds no larger than 4 true leaves. Use the higher rate when the majority of weeds are at the 4 true leaf stage or if henbit, nettle, spurrey or prostrate amaranth are dominant species. |
| Susceptible weeds | Annual poa, black nightshade, clammy goosefoot, henbit, fathen, field pansy, fumitory, groundsel, nettle, prostrate amaranth, redroot, sharp buttercup, shepherds purse, spurrey, willow weed. |
| Water rate | Use a water rate to achieve full coverage of target weeds. |
| No. of applications | No restriction on the number of applications but do not use more 10 L/ha per season. |
| Adjuvants | Do not use any adjuvants with Betanal quattro unless experience has shown them to be crop safe. |
| Compatibility | Betanal quattro is compatible with Lorsban 50EC, Pyramin DF and Versatill Powerflo. Otherwise apply Betanal quattro alone unless experience has shown the mix to be safe. |
| Rainfastness | Avoid treatment if rain is likely before the spray has dried on the foliage. |
| Environmental conditions | Apply Betanal quattro during conditions favouring active weed growth. Avoid application when beet crops are stressed, in drought conditions or when temperatures are above 29°C. |
| Withholding period | Fodder beet, 56 days. Do not graze on treated areas or feed stock with treated crops for 56 days after spraying. Sugar beet, 56 days. Red beet (beetroot), 50 days. |
| Pack size | 5 and 15 litre plastic bottle. Agrecovery compatible. |

Call toll free: 0800 428 246

www.cropscience.bayer.co.nz

Betanal quattro is registered pursuant to the ACVM Act 1997 No. P8851 and is approved pursuant to the HSNO Act 1996, Nos. HSR100882.

Betanal® quattro is a registered trademark of the Bayer Group.

BAYER UTE CARD™ is a trademark of the Bayer Group.

© Bayer CropScience 2015.

For more information on Bayer CropScience products, contact the following Bayer CropScience Territory Managers:

| | | |
|-----------------------|--|-------------|
| Phil Bertram | <i>Upper North Island</i> | 021 426 825 |
| Marc Fox | <i>Gisborne/Hawkes Bay/Nelson/Marlborough</i> | 021 426 823 |
| Jeff Smith | <i>Lower North Island</i> | 021 426 824 |
| David Parker | <i>North and Mid Canterbury</i> | 021 760 794 |
| David Weith | <i>Mid and South Canterbury</i> | 021 426 096 |
| Daniel Suddaby | <i>Otago and Southland</i> | 021 426 822 |
| Colin Dunstan | <i>Seed Treatment Specialist, South Island</i> | 021 323 147 |



Download the new Crop Solutions App from the
Apple App Store or Google Play Store



Science For A Better Life