Confidor®

Net contents : 1, 5, 10 L


Insecticide

A systemic insecticide for the control of thrips in onions, cabbage aphids in vegetable brassicas and aphids in lettuce (cell transplants), and mealybugs on non-bearing grapevines and vines destined for removal.

Contains:350g/litre imidacloprid in the form of a suspension concentrate

Registered pursuant to the ACVM Act 1997, No P5748 See <u>www.foodsafety.govt.nz</u> for registration conditions.

Approved pursuant to the HSNO Act 1996, No HSR002691 See <u>www.epa.govt.nz</u> for approval conditions



UN 3082 PROPER SHIPPING NAME: Environmentally Hazardous Substance, Liquid, N.O.S. (imidacloprid) PACKING GROUP III CLASS HAZCHEM CODE: 3Z

IN A TRANSPORT EMERGENCY DIAL 111 POLICE OR FIRE BRIGADE



WARNING KEEP OUT OF REACH OF CHILDREN HSNO Classifications:



Harmful if swallowed. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects. Very toxic to the soil environment. Very toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates.

PRECAUTIONS:

Read label before use

Do not eat, drink or smoke while using this product.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Avoid contact with the skin and eyes

Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area

Do not apply directly into or onto water

PERSONAL PROTECTION:

When handling, wear full protective clothing such as impervious gloves, respirator, waterproof hat, coat and trousers (worn outside rubber boots).

STORAGE: Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled, move the victim to fresh air immediately.



Confidor is a systemic insecticide of the chloro-nicotinyl group for the control of thrips in onions, cabbage aphids in vegetable brassicas and aphids in lettuce (cell transplants), and mealybugs on non-bearing grapevines and vines destined for removal.

It is an offence to use this product on animals.

DIRECTIONS FOR USE:

Pour Confidor directly into the spray tank with agitators running.

RESISTANCE MANAGEMENT Group 4A Insecticide

Confidor contains a group 4A insecticide and belongs to the neonicotinoid group of chemicals. Resistance to this insecticide could develop from excessive use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies (<u>www.nzpps.org/resistance</u>) that exists for any insect pest on this label.

RECOMMENDATION:

Crop	Pest	Rate	Remarks
Onions	Thrips	225ml/ha	Apply first application when monitoring of thrips levels indicates the threshold level has been reached. Repeat applications at 7- 10 day intervals as a cluster of a maximum of 4 applications. Use in 500-600L of water/ha. The addition of a wetting agent is suggested. Ensure good spray coverage. Refer to the Resistance Management statement.
Vegetable brassicas (cell transplants)	Cabbage aphids	9ml/2.5 litres of water per 1000 plants	Apply evenly in a watering can to seedling trays 1 day before transplanting.
Lettuce (cell transplants)	Aphids	20-30ml/2.5 litres of water per 1000 plants	Apply evenly in a watering can to seedling trays 1 day before transplanting
Grapes – young non- bearing vines and vines for removal	Mealybug	1.5ml/vine in 500ml water per vine	Apply as a single soil drench treatment in autumn or winter around the base of non-bearing replant vines or vines awaiting removal. Ensure the treatment covers the rooting area of the vines. If rainfall or overhead irrigation is not likely in the 2 days after drenching, apply 2L of water per vine once the drench has soaked in.

COMPATIBILITY:

Confidor is compatible with commonly used fungicides



It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD:

Onions	7 days
Vegetable brassicas (cell transplants)	70 days
Lettuce (cell transplants)	35 days

CONDITIONS OF SUPPLY: READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Date of Manufacture: Batch No:

Confidor® is a registered trademark of the Bayer Group

Bayer

Registered to: Bayer New Zealand Limited B:Hive Building, 74 Taharoto Road, Smales Farm, Takapuna, Auckland 0622 Ph. 0800 428 246 EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607



