

Vayego®

Net contents : 1, 2.5, 5, 10, 20 L

Insecticide

A systemic insecticide for the control of codling moth, leafrollers and bronze beetle in apples, pears and nashi and carpophilus beetle in stone fruits.

Contains:


200 g/l tetraniliprole in the form of a suspension concentrate

Registered pursuant to the ACVM Act 1997, No P9623

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, No HSR101424

See www.epa.govt.nz for approval conditions



UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID,
N.O.S. (TETRANILIPROLE SOLUTION)
PACKING GROUP III
CLASS 9
HAZCHEM CODE 3Z

IN A TRANSPORT EMERGENCY DIAL 111 POLICE OR FIRE BRIGADE



28/07/2023

WARNING
KEEP OUT OF REACH OF CHILDREN

GHS Classifications:



Very toxic to aquatic life with long lasting effects
Hazardous to terrestrial invertebrates

PRECAUTIONS:

Read label before use

Do not eat, drink or smoke while using this product.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

Avoid contact with the skin and eyes.

Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

Do not apply directly into or onto water.

Do not apply substance to plants if bees are foraging or the plants are in flower or part flower and are likely to be visited by non-target invertebrate pollinators.

PERSONAL PROTECTION:

When handling, wear full protective clothing such as impervious gloves, respirator, waterproof hat, coat and trousers (worn outside rubber boots).

STORAGE: Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the **PERSONAL PROTECTION** section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum.

Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed, DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin contact occurs, remove contaminated clothing and flush skin with running water. If splashed in eyes, wash out immediately with water, if irritation persists get medical advice. If inhaled, move the victim to fresh air immediately.



28/07/2023

It is an offence to use this product on animals.

DIRECTIONS FOR USE:

MIXING: Half fill the spray tank with water, then pour in the required quantity of **Vayego®** with agitators running, then top up with water.


Do not pre-mix.

RESISTANCE MANAGEMENT:

Group	28	Insecticide
--------------	-----------	--------------------

Vayego® contains tetraniliprole, a diamide insecticide which belongs to the ryanodine receptor modulators mode of action (Group 28). Resistance to this insecticide and related products could develop in some insect pests from repeated use. Use strictly in accordance with label instructions. Avoid treating consecutive insect generations with same mode of action. Successive applications of **Vayego®** or other Group 28 insecticides are acceptable if they are used to treat a single insect generation. Do not rely on **Vayego®** for the control of resistant population. Do not use less than label rates.

RECOMMENDATIONS

Crop	Insect pest	Product Rate	Comments
Apples, pears and nashi	Codling moth, leafrollers and bronze beetle	15 ml/100L as a dilute application	Make up to 2 applications per season. The first application should be made post-petal fall. A second application may be made as required, with a minimum of a 21 day interval and up to 14 days before harvest, as part of an IPM program. The second application should only be made if pest thresholds are exceeded. NB: Do not apply more than two consecutive applications of Group 28 Insecticides in any one season. For dilute application, spray to the point of run-off. Ensure complete coverage. For concentrate spraying adjust the rate accordingly.
Stone fruit	Carpophilus beetle 	15 mL/100 L	Monitor stone fruit orchards for beetles as fruit approaches maturity and become susceptible to attack. Commence application before beetle populations reach damaging levels and re-apply treatments if necessary, at an interval of 14-21 days. If significant rainfall is predicted within 24 hours of application, ensure that the crop is dry at application and sufficient drying time is allowed prior to rainfall. Do not apply more than two applications per season in each crop. Apply in sufficient water, to the point of runoff. No adjuvant is necessary.

PHYSICAL COMPATIBILITY

Vayego® is compatible with commonly used crop protection products, contact your local Bayer representative for further details.

RESTRICTIONS

Pome fruit: maximum 2 applications of 60 g ai/ha per year with a minimum of 21-day interval between applications.

Stone fruit: maximum 2 applications of 60 g ai/ha per year with a minimum of 14-day interval between applications.

Vayego must not be applied when wind speeds are less than 3 km/hr or more than 20 km/hr as measured at the application site.

Vayego can only be applied by ground-based methods.

Vayego can only be applied post-bloom

When applied on pome fruit and stone fruit, Vayego must not be applied within 25 m of any waterbody.

Use best available application technique to minimise off-target drift.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS:

Re-entry intervals: Do not enter treated area before the spray deposit has dried completely (for at least 12 hours after the application)

Apples, pears and nashi: 14 days before harvest.
Import tolerances for produce treated with Vayego may be pending in some countries. Consult with your exporter or Bayer NZ before applying Vayego to export crops.

Stone fruits: 4 days before harvest.
Import tolerances for produce treated with Vayego may be pending in some countries. Consult with your exporter or Bayer NZ before applying Vayego to export crops.



28/07/2023

CONDITIONS OF SUPPLY: READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Batch No:

Vayego® is a registered trademark of the Bayer Group

Bayer

Registered to:
Bayer New Zealand Limited
B:Hive Building
74 Tahoroto Road
Takapuna
Auckland 0622

Ph. 0800 428 246
EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607

A handwritten signature in blue ink, appearing to be "John Zimmo".

28/07/2023