

TOXIC
Keep out of reach of children

Mesurool[®] 200 SC

Contains: 200g/litre methiocarb in the form of a suspension concentrate.

Net Contents: 1 litre / 3 litre / 5 litre

Bird Repellent

For repelling birds from crops.

Approved pursuant to the HSNO Act 1996, No HSR100895
See www.epa.govt.nz for approval conditions

APPROVED HANDLER: This product must be under control of an approved handler when applied in a wide dispersive manner or when used by a commercial contractor.

HSNO Classifications: 6.1D, 6.9B, 9.1A, 9.2A, 9.3A, 9.4A



WARNING:

Toxic – may be harmful if swallowed.

Harmful-presumed to/may cause organ damage from repeated oral exposure at high doses.

Very toxic to aquatic organisms – must not be applied directly onto, into or over water.

Very toxic to the soil environment.

Very toxic to terrestrial vertebrates.

Very toxic to terrestrial invertebrates

PRECAUTIONS:

PERSONAL PROTECTION: Do not eat, drink or smoke while using. May be harmful if swallowed, inhaled or absorbed through the skin. Material treated with this substance must not be used for human or animal consumption.

Wear protective clothing such as impervious gloves, goggles, waterproof hat, coat and trousers when using. Appropriate personal protection equipment shall be used when handling material treated with this substance. Remove protective clothing and wash hands and face thoroughly before meals and after work.

STORAGE: Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from foodstuffs. Treated material must be completely covered by soil when sown. Excess treated material must not be left in areas accessible to birds.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

Disposal: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container. Material treated with this substance shall be disposed of in accordance with the disposal requirements for toxic and ecotoxic substances

Toxic to bees: Spray must not contact plants in flower if they are likely to be visited by bees.

Container Disposal: Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

FIRST AID:

If swallowed do NOT induce vomiting.

For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

If skin contact occurs, remove contaminated clothing and flush skin with running water, if irritation occurs seek medical advice.

If splashed in eyes, wash out immediately with water.

If inhaled move the victim to fresh air immediately. Begin artificial respiration if the victim is not breathing. Use mouth-to-nose rather than mouth-to-mouth. Obtain medical attention.

Mesurol 200 SC is a bird repellent. It is very effective for repelling birds which can damage seeds and seedlings and flower buds of kiwifruit.

Mixing:

Pour **Mesurol 200 SC** directly into the spray tank with agitators running.

COMPATIBILITY:

Mesurol 200 SC is compatible with the most commonly used fungicides, insecticides and herbicides. Do not use with alkaline materials such as Bordeaux mixture and lime sulphur.

TO REPEL BIRDS:

Crop	Rate	Comments
Seedlings: Broccoli Brussels sprouts Cabbage Cauliflower Celery Lettuce Tomatoes	3.75 L/ha as a band spray over the crop row	Make the first application immediately after the crop emerges and repeat if necessary 14 days later so long as the crop is still vulnerable to damage by birds.
Peas	1.5 L/100kg of seed	Add the required quantity of Mesurol 200 SC and mix thoroughly to ensure complete coverage. A concrete mixer is satisfactory. When sowing peas use Mesurol 200 SC treated seed for the two outside runs. After this it may be sufficient to sow treated seed from one of the drill's four hoppers.
Kiwifruit	375 ml/100 litres and apply to the point of run-off using 800-1000 litres of water/ha.	Apply during August and September at the first sign of bird damage, or as soon as birds are present in the orchard in numbers, and buds have reached a susceptible stage. A single application is generally sufficient, unless heavy rain occurs when a re-application may be necessary.

BEE SAFETY:

MESUROL 200 SC IS TOXIC TO BEES. Follow best practice when spraying:

1. Do not apply to any flowering plants.
2. Do not apply to shelter belts.
3. Mow sward prior to applying to remove any flowering weeds which may attract bees.
4. Only spray when bees are not actively foraging, before 8.00 am or in the late afternoon/early evening.

WITHHOLDING PERIOD:

Vegetable crops21 days

Kiwifruit.....Do not apply after start of flowering

CONDITIONS OF SUPPLY:

READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Mesurool® is a registered trademark of the Bayer Group

Date of Manufacture:

Batch No:

Bayer

Bayer New Zealand Limited
3 Argus Place, Hillcrest, Auckland 0627
Ph. 0800 428 246
EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607

