

# Galmano®

**Contains: 167g/litre fluquinconazole in the form of a flowable suspension concentrate**

**Net Contents:  
5/10/ 200 litres**

## ***Fungicide***

Seed treatment fungicide for control of foliar diseases in wheat.

Registered pursuant to the ACVM Act 1997, No P7795  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to the HSNO Act 1996, No HSR100402  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

  
Marco D'Amico  
Manager Approvals Operations  
Regulation and Assurance



08 FEB 2017

**Approved Handler:** This product must be under control of an approved handler during use.

**Tracking:** A record must be kept of the use and disposal of this product.

**HSNO Classifications:** 6.1D, 6.3A, 6.4A, 6.5B, 6.8C, 6.9A, 9.1A, 9.3C



**DANGER: Keep out of reach of children**

Harmful – may be harmful if swallowed, inhaled or absorbed through the skin.

Harmful – may cause skin irritation.

Harmful – may cause eye irritation.

Harmful – may cause sensitisation from prolonged skin contact.

Harmful - may cause reproductive/developmental damage from repeated oral exposure at high doses.

Toxic – may cause target organ damage from repeated oral exposure at high doses.

Very toxic to aquatic organisms with long-lasting effects.

Harmful to terrestrial vertebrates.

**PRECAUTIONS:**

**PERSONAL PROTECTION:**

Do not eat, drink or smoke while using.

Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using.

Remove protective clothing and wash hands and face thoroughly before meals and after work.

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation. Avoid contact with the eyes.

**STORAGE:** Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs.

**SPILLS AND DISPOSAL:** When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section.

Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

**DISPOSAL:** Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container.

**CONTAINER DISPOSAL:** Triple rinse container and add residue to mixture. Take to an Agrecovery collection site.

**FIRST AID:**

If swallowed do NOT induce vomiting.

For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

If skin contact occurs, remove contaminated clothing and flush skin with running water.

If splashed in eyes, wash out immediately with water.

If inhaled move the victim to fresh air immediately.



**Containers of treated seed must be labelled as follows:**

1. **Seed treated with Galmano must not be used for animal or human consumption.**
2. **Treated seed must be completely covered by soil when sown.**
3. **Excess treated seed must not be left in areas accessible to birds.**
4. **It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**
5. **Seeds treated with this substance shall be disposed of in accordance with the disposal requirements for toxic and ecotoxic substances.**
6. **Withholding period: Stock must not graze wheat within 42 days of planting.**

**[These statements must be in bold print on the labels of treated seed containers]**

**Galmano** is a systemic triazole seed treatment fungicide for the control of powdery mildew, leaf rust and stripe rust in wheat.

**It is an offence to use this product on animals.**

**RECOMMENDATIONS:**

<b>Crop</b>	<b>Disease</b>	<b>Remarks</b>
Wheat	Powdery mildew Leaf rust Stripe rust	Mix 1.0 litre with sufficient water (5-10 litres) and apply as a slurry treatment to 1.0 tonne of seed ensuring complete coverage.

**Application Method:**

**Galmano** may be applied through slurry, Mistomatic, Sutton and similar seed treatment machinery.

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds**

**Withholding period:**

Wheat.....42 days from planting



08 FEB 2017

**CONDITIONS OF SUPPLY:**

**READ THESE CONDITIONS OF SUPPLY BEFORE BUYING AND USING. IF THESE CONDITIONS ARE UNACCEPTABLE, RETURN AT ONCE UNOPENED.**

To the extent permitted by law Bayer does not accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with these goods, including without limitation their storage, application, handling or use. Bayer does not make any warranties (express or implied) with respect to fitness for purpose, merchantable quality or any other matter.

To the extent only that Bayer is prevented by law from contracting out of the Consumer Guarantees Act, then these conditions apply subject to that Act. The Act shall not apply where these goods are acquired for business purposes.

Date of Manufacture:

Batch No:

Galmano® is a registered trademark of the Bayer Group

Bayer

**Registered to:**

Bayer New Zealand Limited

3 Argus Place, Hillcrest, Auckland 0627

and Christchurch

Ph. 0800 428 246

EMERGENCY RESPONSE SERVICE (24 hrs) 0800 734 607



Merce Zindley  
Manager Approvals Operations  
Regulation and Assurance



08 FEB 2017